Estuaries Vol. 13, No. 4, p. 513–515 December 1990

Subject Index, Volume 13

abundance model, 486
Achirus mazatlanus, 219
acoustic scatter, 1
Advanced Very High Resolution Radiometer, 155
aerial imagery, 272*
Alaskan fjords, 380
alkalinity, 8
aluminosilicate, 8
aluminum, 8, 227*
Amazon River, 204
Ammodytes sp., 43
ammonia, 404, 418
ammonium, 219*
amphipod(s), 174, 301*, 332*, 431
Argopectens irradians concentricus, 322*
Arius liropus, 219
arsenic, 227*
assimilation, 20
astronomical tides, 155
Atlantic croaker, 453
Atlantic menhaden, 453
Avicennia, 204
Avicennia bicolor, 182
Avicennia germinans, 219

bar-built estuaries, 504
Barataria Basin, Louisiana, 393, 404
barotropic exchange, 258*
bay scallop, 322*
Bay St. Louis, Mississippi, 332*
Beaufort Inlet, North Carolina, 453
benthic community, 219*
benthic predators, 462
biomass, 174
Biscayne Bay, Florida, 227*
black mangrove, 204
boat-generated pollution, 282*
Boca de Quadra, Alaska, 380
Bogue Sound, North Carolina, 322*
bongo nets, 453
bottom-water intrusions, 31
box model, 418
brackish community, 373
brackish marsh(es), 332*, 404
Brazilian Coast, 204
Brevoortia patronus, 14
Brevoortia tyrannus, 14, 453

cadmium, 8, 227*
Calcasieu Lake, Louisiana, 258*
Callinectes sapidus, 51, 431
canals, 272*
Cananeia Lagoon, 193
Cancer magister, 59, 469
Capitella cf. *capitata*, 174
carbonates, 161
Catostomus, 43
cellulase activity, 14
Central America, 182
Centropomus robalito, 219
cesium-137, 380
Charleston Harbor, South Carolina, 1
Chesapeake Bay, 236*, 282*, 287*, 431, 486
chlorophyll *a*, 441
chromium, 227*
Chrysaora quinquecirrha, 486
circulation features, 31
circulation model, 236*
Cladophora, 479
clay mineral composition, 161
climatic seasonality, 204, 219
climatological variations, 265*
coastal plain estuaries, 504
cobalt, 8
community structure, 469
Conocarpus, 204
contaminants, 81
contamination by metals, 227*
Cooper River, South Carolina, 373
copper, 8, 227*
Core Sound, North Carolina, 322*
Corophium louisianum, 332*
corrosion, 89
crabs, 469
Crangon franciscorum, 469
Crangon nigricauda, 469
cresote, 72
current(s), 31, 133, 258*
current measurements, 250*
current speed, 250*
current velocity, 118
Cyclotella caspia, 287*
Cyclotella meneghiania, 287*
Cyclotella striata, 287*
Cymadusa compta, 301*
Cymatogaster aggregata, 469
Cyprinodon variegatus, 20

Dasybranchus lumbricoides, 174
decomposition, 193
degradation, 219
deltaic coast, 272*
deltaic cycle, 404
demersal fish, 469
density currents, 125
density gradient, 1, 118
density stratification, 349
destratification, 236*
detritivores, 20
detritus, 14
detritus feeding, 20
Diapterus peruvianus, 219
diatoms, 287*
2,4-dichlorophenoxyacetic acid (2,4-D), 72
diffusion, 81
diptera, 341*
discriminant analysis, 287*
dispersion, 133
dispersion processes, 118
dissolved nutrients, 404, 418
dissolved organic carbon, 404
dissolved organic nitrogen, 404
dissolved oxygen, 8, 236*
dissolved silicon, 418
diurnal tides, 362
diversity of estuaries, 504
dolochopodid, 341*
drilling fluid, 89
drowned river valleys, 504
dry weather mangroves, 182
Dungeness crab, 59

ebb-tidal fronts, 1
Elbe Estuary, 161
elevations, 133
English sole, 59, 469
Enophrys bison, 469
entrance sill, 31
epibenthic crustaceans, 469
episodic deposition, 380
Erichsonella attenuata, 431
estuaries, definition, 492
Eucinostomus, 219
Eurytium limosum, 462
evapotranspiration, 204
export, 418
external inputs, 418

feeding habits, 462
fiddler crabs, 332*
fish, 193
fish community, 219
fish diet, 332*
fish larvae, 453
fishery science, 231
fjord, 31, 504
flumes, 393
flushing times, 133
flux(es), 118, 404, 418, 404
food sources, 14
forecasting, 486
forest, 182
Fourleague Bay, Louisiana, 404
frequency-dependent model, 362
freshwater, 145
freshwater discharges, 373
freshwater input(s), 125, 404
freshwater lens, 1
freshwater tidal river, 441
Fundulus grandis, 332*
Fundulus heteroclitus, 20, 462

* Vol. 13(3)—These numbers overlap with Vol. 13(2).

© 1990 Estuarine Research Federation

0160-8347/90/040513-03$01.50/0

Gadus morhua, 43
Galeichthys caerulescens, 219
Gammarus mucronatus, 431
gastropoda, 479
Geum Estuary, Korea, 8
grain size fractions, 161
Grandidierella bonnieroides, 174
Grays Harbor, Washington, 59
Great Lakes estuaries, 493, 508
Great Sippewissett, 20
Gulf killifish, 332*
Gulf menhaden, 14
Gulf of Mexico, 89, 98, 258*, 272*

Halodule wrightii, 51
Hargeria rapax, 332*
Hippoglossoides platessoides, 43
Hudson Estuary, 362
humic substances, 193
Hydrilla verticillata, 311*
Hydrobia truncata, 479
hydrobiid snails, 479
hydrocarbons, 89, 98
hydrologic changes, 272*

ichthyoplankton, 43
impounded salt marsh, 25
inlets, 504
intertidal distribution, 462
intra-tidal variation, 118
iron, 8
isopod, 431
isotopes, 161

James River, 287*
Juan de Fuca estuary, 31

killifish, 462
Kjeldahl nitrogen, 98, 404

laboratory simulation, 81
lacustrine estuary, 506
Lagodon rhomboides, 51, 453
Laguna Joyuda, 174
Laguncularia, 193, 204
Laguncularia racemosa, 219
Lake Erie, 493
Lake Ontario, 493
Lake Superior, 493
larval flies, 341*
lead, 8, 227*
lead-210, 380
Leiostomus xanthurus, 453
Leptocottus armatus, 469
Leptocylindrus minimus, 287*
light attenuation, 441
light conditions, 311*
Lile stolifera, 219
Liparis sp., 43
Littoridinops palustris, 332*
Liverpool Bay, 125
Long Island Sound, 362
long-term trends, 265*
longitudinal dispersion, 118
Louisiana, 72
Louisiana estuaries, 265*
Louisiana wetlands, 272*
Lumpenus sagitta, 469

macrobenthic, 174
macroinfaunal community, 174
macrophytes, 311*
macrotidal estuary, 133
Mallotus villosus, 43
management, 231
manganese, 8
mangrove(s), 182, 193, 204
mangrove ecology, 219
Manicouagan River, 43
marina facilities, 282*
marine fishes, 453
marsh plants, 265*
marsh subsidence, 337*
marsh-water column interactions, 393, 404
medusal counts, 486
mercury, 227*
mesohaline, 287*, 441
metals, 89, 227*, 301*
meteorologic forcing, 393
Mexican Pacific, 219
Miami River, Florida, 227*
microalgal community, 219*
microcosm(s), 301*, 311*
Micropogonias undulatus, 453
migration, 59
Mississippi Sound, 98
Mobile Bay, 145, 155
modeling, 236*
models, 258*
Montrichardia, 204
mud crab, 462
Mugil curema, 219
muskrats, 337*
Myriophyllum spicatum, 311*
Myrophis punctatus, 453

^{15}N assimilation, 20
napthalene, 72, 81
Narragansett Bay, Rhode Island, 418
natural tracers, 161
Neomysis mercedis, 469
New England, 25
Newport River, North Carolina, 453
nickel, 8, 227*
nitrate, 219*, 404, 418
nitrogen, 20, 287*
nonanthropogenic metals, 161
North Sea, 161
Nototropis sp., 43
numerical hydrodynamical model, 349
nutrient(s), 193, 418
nutrient concentrations, 418
nutrient dynamics, 404
nutrient exchange, 219*

$^{16}O/^{18}O$ ratio, 161
oceanic tides, 362
octadecane, 72
oil drilling activities, 81
Old Woman Creek estuary, 493
oligochaete, 174
oligohaline, 287*, 341*
Ondatra zibethicus, 337*
organic carbon, 98
organic matter, 161
Osmerus mordax, 43
Outer Banks, North Carolina, 322*

p-dichlorobenzene, 81
Pacific Coast, 182
Palaemonetes intermedius, 51
Parker River National Wildlife Refuge, Massachusetts, 479
Parophrys vetulus, 59, 469
particle transport, 161
particulate carbon, 418
particulate matter, 193
particulate nitrogen, 418
particulate transport, 118
partly-mixed estuary, 349
Pelastoneurus abbreviatus, 341*
Penaeus duorarum, 51
pentachlorophenol (PCP), 72
periphyton, 193
petroleum drilling, 89
petroleum hydrocarbons, 72
Phanerodon furcatus, 469
phenols, 98
phosphate, 219*, 418
phosphorus, 287*
photosynthetic production, 193
Phragmites australis, 25
phytoplankton, 1, 14, 193, 287*
pinfish, 453
plant communities, 373
plant density, 182
Platichthys stellatus, 469
point-source inputs, 418
pollutant profiles, 98
polychaetes, 51
polychlorobiphenyls (PCBs), 72
Port Townsend, 31
portunid decapod, 431
Potamogeton perfoliatus, 311*
Potomac River, 441
primary production, 193
primary productivity, 219
production, 479
Providence River, 418
Pseudopleuronectes americanus, 43
Pterocarpus, 204
Ptychodiscus brevis, 322*
Puerto Rico, 174
Puget Sound, 31
pump-out facilities, 282*

radiometric measurments, 441
Rappahannock River, 287*
recreational boaters, 282*
recruitment failure, 322*
red tide, 322*
rediversion, 373
redox conditions, 72
remote sensing, 155
residual currents, 349
restoration, 25
Rhacochilus vacca, 469
Rhizophora, 193, 204
Rhizophora mangle, 219
river discharge, 155, 219
river-dominated estuary, 145
riverine input, 1

runoff, 133
Ruppia maritima, 431

St. Lawrence Estuary, 43
St. Lawrence River, 493
salinity, 31, 118, 133, 145, 236*, 311*
salinity change(s), 373
salinity gradient, 1
salinity trends, 265*
salt balance, 133
salt marsh killifish, 20
salt marsh pools, 479
salt wedge, 161
Sao Paulo, Brazil, 193
Sapelo Island, Georgia, 462
Scomber scombrus, 43
sea level variability, 362
sea nettle, 486
seagrass, 301*, 431
seagrass beds, 51
seasonal infestation, 486
Sebastes fasciatus, 43
Secchi depth, 441
secondary production, 431, 479
sediment, 72, 89
sediment accumulation, 380
sediment bioassays, 98
sediment compaction, 272*
sediment dynamics, 393
sediment exchanges, 393
sediment profiles, 380
sediment transport, 380
sedimentary contaminants, 98
sedimentation, 380
sediments, 81, 161, 193, 219*, 227*
Seekonk River, 418
semidiurnal tides, 362
sewage treatment plants, 418
shear-induced dispersion, 118
Shot-Sensor method, 250*
shrimp, 469
Skeletonema costatum, 287*
Skeletonema potamos, 287*
sludge, 89
Smeaton Bay, Alaska, 380
Spartina, 14, 204
Spartina alterniflora, 20, 25, 373
Spartina cynosuroides, 341*
spawning, 453
speckled worm eel, 453
spoil banks, 272*
spot, 453
Spurwinkia salsa, 479
stable isotopes, 14
Steninonereis martini, 174
stirring, 125
Stonington, Connecticut, 25
storms, 236*
stratification, 125, 145
stream flow, 486
submersed macrophytes, 441
subsiding marshes, 404
subtidal exchange, 258*
subtidal frequencies, 362
suspended matter, 161
suspended particulate material, 441
suspended sediment, 155
suspended solids, 118, 393

Tamar Estuary, England, 118, 133, 349
temperature patterns, 155
temporal abundances, 453
temporal variability, 155
Thalassia, 301*
Thalassodrilides gurwitschi, 174
Thinophilus frontalis, 341*
tidal flooding, 337*
tidal frquency, 362
tidal prism mixing model, 373
tidal range, 441
tidal straining, 125
tides, 362
time scales, 145
total nitrogen, 404
total organic carbon (TOC), 89, 98
toxic organics, 72
toxicity, 98
trace metals, 8
transparency, 193
transport, 161
trawl surveys, 59
tributylin, 301*
1,1,1-trichloro-2,2-bis(4-chlorophenyl) (DDT), 72
tropical lagoon, 174
turbidity gradient, 1
Typha angustifolia, 25

Uca longisignalis, 332*

Vallisneria americana, 311*
vertical current profiles, 349
vertical eddy viscosity, 349
vertical shear, 118

water pollution, 282*
water quality, 282*
wetland fragmentation, 272*
wetland loss, 265*, 272*
wildlife, 337*
Willapa Bay, Washington, 59
wind mixing, 236*
wind-driven model, 258*

xanthis crab, 462

Yaquina Bay, Oregon, 469
York River, 287*
York River, Virginia 219*

zinc, 8, 227*
zooplankton, 1, 14, 193
Zostera marina, 431